









**Spring 2018 Organic II Final Exam****100pts (graded as 150pts)****Name:**

If you do not want your graded exam placed in the box outside my office, then please check here

1) Identify the class of compounds (functional group) each of the following molecules belongs to. (10pts)



2) Put a cross through the molecule (above) that is unstable above room temperature and will liberate nitrogen gas. (1pt)

3) Define the following terms (5x2=10pts)

(a) *Carboxylic Acid Derivative*

(b) *Thermodynamic Control*


(c) *Condensation Reaction*

(d) *Acyl Group*

(e) *Hammond's Postulate*


4) (1+4=5pts) Provide one of the factors that influence the relative rates of Nucleophilic Acyl Substitution reactions for carboxylic acid derivatives (e.g. *esters* are more reactive than *amides*).

Draw the mechanism (*i.e. curly arrows*) for this reaction which occurs under *basic* conditions.











5) Write the mechanism (*i.e. curly arrows*) for the **acid** catalyzed nucleophilic addition of water to acetone (propanone), producing the ketone hydrate. (5pts)

6) (1+2+2+2=7pts) The following molecule was produced in a [4+2] cycloaddition (Diels-Alder) reaction.






- a) The right hand side ring contains a cyclic version of what type of functional group ?
- b) Draw the *diene* and *dienophile* which would react together to give this product, and then draw the **mechanism** (*i.e. curly arrows*) for this reaction.
- c) If the above molecule was hydrolyzed using water, draw the product dicarboxylic acid.









7) Indicate which of the following molecules are *aromatic*, *non-aromatic* or *anti-aromatic*. (Assume all the molecules are planar). (8pts)



8) Circle the *strongest acid* (proton donor) in the following threesomes. (3pts)






9) Fill in the blanks (either SM/products) for six of the following reactions. (6x2=12pts)




10) Give the products in **four** of the following transformations. (8pts)






11) Circle the *most basic atom* in each of the following species. (3pts)



12) Give reagents for the following transformations. (4x3=12pts)



13) Name two of the following compounds in IUPAC form (6pts).




14) Draw in line angle (*stick figure*) form, two of the following named molecules. (6pts)

*2-Bromo-3-hydroxybutanoic acid lactone*

*N, N-Diethylpropanamide*

*2-Bromopropanoyl Bromide*

15) Write the mechanism (*i.e. curly arrows*) for this generic Electrophilic Aromatic Substitution (EAS reaction) of Benzene. (4pts)



**\*\*\*Bonus question\*\*\*** (up to 3 points)

Draw the structure of Pyridine ( $C_5H_5N$ ), and using the Polygon Rule, *justify* that it has an aromatic  $\pi$  system.

hydrogen	<b>H</b>	1.0079	1	helium	<b>He</b>	4.0026
lithium	<b>Li</b>	6.941	3	boron	<b>B</b>	10.811
beryllium	<b>Be</b>	9.0122	4	carbon	<b>C</b>	12.011
sodium	<b>Na</b>	22.990	11	nitrogen	<b>N</b>	14.007
magnesium	<b>Mg</b>	24.305	12	oxygen	<b>O</b>	15.999
potassium	<b>K</b>	39.098	19	fluorine	<b>F</b>	18.998
calcium	<b>Ca</b>	40.078	20	neon	<b>Ne</b>	20.180
scandium	<b>Sc</b>	44.956	21	boron	<b>B</b>	26.982
yttrium	<b>Ti</b>	47.867	22	carbon	<b>C</b>	31
strontium	<b>V</b>	50.942	23	nitrogen	<b>N</b>	30
ruthenium	<b>Cr</b>	51.996	24	oxygen	<b>O</b>	30
ytterbium	<b>Mn</b>	54.938	25	fluorine	<b>F</b>	30
zirconium	<b>Fe</b>	55.945	26	neon	<b>Ne</b>	30
cerium	<b>Co</b>	58.933	27	boron	<b>B</b>	35
europium	<b>Ni</b>	58.993	28	carbon	<b>C</b>	35
gadolinium	<b>Cu</b>	63.946	29	nitrogen	<b>N</b>	35
thulium	<b>Zn</b>	68.939	30	oxygen	<b>O</b>	35
lanthanum	<b>Ga</b>	69.723	31	fluorine	<b>F</b>	35
neodymium	<b>Ge</b>	72.61	32	neon	<b>Ne</b>	36
praseodymium	<b>As</b>	74.922	33	boron	<b>B</b>	36
holmium	<b>Se</b>	78.96	34	carbon	<b>C</b>	36
dysprosium	<b>Br</b>	79.941	35	nitrogen	<b>N</b>	36
erbium	<b>Kr</b>	83.80	36	oxygen	<b>O</b>	36
thulium	<b>Al</b>	86.948	37	fluorine	<b>F</b>	36
ytterbium	<b>Si</b>	90.948	38	neon	<b>Ne</b>	36
lutetium	<b>P</b>	92.948	39	boron	<b>B</b>	36
cerium	<b>S</b>	93.948	40	carbon	<b>C</b>	36
europium	<b>Cl</b>	95.948	41	nitrogen	<b>N</b>	36
thulium	<b>Ar</b>	98.948	42	oxygen	<b>O</b>	36
ytterbium	<b>Fr</b>	101.948	43	fluorine	<b>F</b>	36
thulium	<b>Rb</b>	102.948	44	neon	<b>Ne</b>	36
ytterbium	<b>Sr</b>	102.948	45	boron	<b>B</b>	36
ytterbium	<b>Zr</b>	102.948	46	carbon	<b>C</b>	36
ytterbium	<b>Nb</b>	102.948	47	nitrogen	<b>N</b>	36
ytterbium	<b>Mo</b>	102.948	48	oxygen	<b>O</b>	36
ytterbium	<b>Tc</b>	102.948	49	fluorine	<b>F</b>	36
ytterbium	<b>Ru</b>	102.948	50	neon	<b>Ne</b>	36
ytterbium	<b>Rh</b>	102.948	51	boron	<b>B</b>	36
ytterbium	<b>Pd</b>	102.948	52	carbon	<b>C</b>	36
ytterbium	<b>Ag</b>	102.948	53	nitrogen	<b>N</b>	36
ytterbium	<b>Cd</b>	102.948	54	oxygen	<b>O</b>	36
ytterbium	<b>In</b>	102.948	55	fluorine	<b>F</b>	36
ytterbium	<b>Sn</b>	102.948	56	neon	<b>Ne</b>	36
ytterbium	<b>Sb</b>	102.948	57	boron	<b>B</b>	36
ytterbium	<b>Te</b>	102.948	58	carbon	<b>C</b>	36
ytterbium	<b>I</b>	102.948	59	nitrogen	<b>N</b>	36
ytterbium	<b>Xe</b>	102.948	60	oxygen	<b>O</b>	36
ytterbium	<b>Fr</b>	102.948	61	fluorine	<b>F</b>	36
ytterbium	<b>Ra</b>	102.948	62	neon	<b>Ne</b>	36
ytterbium	<b>Lr</b>	102.948	63	boron	<b>B</b>	36
ytterbium	<b>Rf</b>	102.948	64	carbon	<b>C</b>	36
ytterbium	<b>Db</b>	102.948	65	nitrogen	<b>N</b>	36
ytterbium	<b>Sg</b>	102.948	66	oxygen	<b>O</b>	36
ytterbium	<b>Bh</b>	102.948	67	fluorine	<b>F</b>	36
ytterbium	<b>Hs</b>	102.948	68	neon	<b>Ne</b>	36
ytterbium	<b>Mt</b>	102.948	69	boron	<b>B</b>	36
ytterbium	<b>Uuu</b>	102.948	70	carbon	<b>C</b>	36
ytterbium	<b>Uub</b>	102.948	71	nitrogen	<b>N</b>	36
ytterbium	<b>Uog</b>	102.948	72	oxygen	<b>O</b>	36

\* Lanthanide series

lanthanum	cerium	praseodymium	neodymium	europium	gadolinium	terbium	dysprosium	yttrium	thulium	ytterbium	ytterbium
57	58	59	60	61	62	63	64	65	66	67	68
138.91	140.12	140.91	144.24	145.93	149.36	151.96	153.93	155.93	157.93	159.93	161.93
La	Ce	Pr	Nd	Pm	Sm	Eu	Gd	Tb	Dy	Ho	Er
cerium	praseodymium	neodymium	europium	gadolinium	terbium	dysprosium	yttrium	thulium	ytterbium	ytterbium	ytterbium
89	90	91	92	93	94	95	96	97	98	99	100
140.91	141.91	142.91	143.91	144.91	145.91	146.91	147.91	148.91	149.91	150.91	151.91
Ac	Th	Pa	U	Np	Pu	Am	Cm	Bk	Cf	Es	Fm
actinium	thulium	protactinium	uranium	nepalium	plutonium	curium	berkelium	californium	einsteiniun	fermium	moscovium
122	232.04	231.04	233.03	234.03	235.03	236.03	237.03	238.03	239.03	240.03	241.03

\* \* Actinide series









## Spring 2018 Organic II Final Exam

100pts (graded as 150pts)

Name: HIYAM DUNN

*If you do **not** want your graded exam placed in the box outside my office, then please check here* 

1) Identify the class of compounds (functional group) each of the following molecules belongs to. (10pts)



2) Put a cross through the molecule (above) that is unstable above room temperature and will liberate nitrogen gas. (1pt)

3) Define the following terms (5x2=10pts)

(a) *Carboxylic Acid Derivative* A functional group that can be hydrolyzed to a carboxylic acid.

(b) *Thermodynamic Control* When a reaction produces the most stable product.

(c) *Condensation Reaction* A reaction where two (or more) molecules add together along with the expulsion of a small molecule.


(d) *Acyl Group* Part of a molecule that has a C=O double bond.

(e) *Hammond's Postulate* For related processes, things that are similar in energy will be similar in structure.


4) (1+4=5pts) Provide **one** of the factors that influence the relative rates of Nucleophilic Acyl Substitution reactions for carboxylic acid derivatives (e.g. esters are more reactive than amides).

- Leaving group ability
- Amount of resonance in the SN.


Draw the mechanism (*i.e. curly arrows*) for this reaction which occurs under *basic* conditions.



5) Write the mechanism (*i.e. curly arrows*) for the **acid** catalyzed nucleophilic addition of water to acetone (propanone), producing the ketone hydrate. (5pts)




6) (1+2+2+2=7pts) The following molecule was produced in a [4+2] cycloaddition (Diels-Alder) reaction.




a) The right hand side ring contains a cyclic version of what type of functional group?

Anhydride

b) Draw the *diene* and *dienophile* which would react together to give this product, and then draw the **mechanism** (*i.e. curly arrows*) for this reaction.




c) If the above molecule was hydrolyzed using water, draw the product dicarboxylic acid.




7) Indicate which of the following molecules are *aromatic*, *non-aromatic* or *anti-aromatic*. (Assume all the molecules are planar). (8pts)




Non




Non




Aromatic




Non




Non



Aromatic






Non





Non

8) Circle the *strongest acid* (proton donor) in the following threesomes. (3pts)




(a)









(b)








(c)






9) Fill in the blanks (either SM/products) for six of the following reactions. (6x2=12pts)




10) Give the products in **four** of the following transformations. (8pts)



11) Circle the *most basic atom* in each of the following species. (3pts)




12) Give reagents for the following transformations. (4x3=12pts)




13) Name two of the following compounds in IUPAC form (6pts).



Propionic Ethanoic  
Anhydride




Ethyl-2-Methyl  
Propanoate




But-3-enal

14) Draw in line angle (*stick figure*) form, two of the following named molecules. (6pts)


### *2-Bromo-3-hydroxybutanoic acid lactone*




### *N,N-Diethylpropanamide*



### *2-Bromopropanoyl Bromide*



15) Write the mechanism (*i.e. curly arrows*) for this generic Electrophilic Aromatic Substitution (EAS reaction) of Benzene. (4pts)



\*\*\*Bonus question\*\*\* (up to 3 points)

Draw the structure of Pyridine ( $C_5H_5N$ ), and using the Polygon Rule, *justify* that it has an aromatic  $\pi$  system.

